



Brian B. Gilmer, MD US Ski Team Physician

Karly M. Dawson PA-C Teaching Associate

REHABILITATION GUIDELINES FOR MICROFRACTURE/BIOCARTILAGE OF FEMORAL CONDYLE

PHASE I (0-6 WEEKS) DATES:	
Appointments	 Begin physical therapy 3 days post-op 1 – 2 x per week
Rehabilitation Goals	 Protection of the post-surgical knee Restore normal knee range of motion and patellar mobility Eliminate effusion Restore leg control
Precautions	 Femoral Condyle: NWB x 6 weeks Brace locked in extension x 1 week, then unlock 0-90 degrees in weight bearing x 6 weeks Progress ROM as tolerate no restrictions Patellofemoral: Avoid open chain strength Weight bearing 0-90 degrees closed kinetic chain strength with brace
Suggested Therapeutic Exercises	 Flexibility exercises 0-2 weeks: Quad sets (not w/PF), SLR, calf pumps, passive leg hangs to 90 2-6 week: Quad, hamstring, and glut sets, SLR, side-lying hip and core
Suggested Cardiovascular Exercise	Upper body circuit training or upper body ergometer
Progression Criteria	 6 weeks post-op No effusion Full knee extension

PHASE II (6-8 WEEKS)	DATES:
Appointments	Continue PT 2x week
Rehabilitation Goals	 Full ROM – progress as tolerated Improve gait mechanics Progress lower extremity strength and endurance Gradually increase functional activities
Precautions	 Advance weight bearing 50% weekly until full at 8 weeks post-op Avoid loading knee at deep flexion angles, no squat > 90 degrees
Suggested Therapeutic Exercises	Advance Phase I exercises, gait drills, hip and core strengthening; Begin weight bearing exercises, weight shifts, calf raise, TKE with T-Band, Bridging, low resistance squats.
Progression Criteria	 Full ROM Good tolerance to weight bearing progression No effusion Normal gait pattern

PHASE III (8-12 WEEKS)	DATES:
Appointments	Continue PT 1-2x week
Rehabilitation Goals	 Restore normal gait mechanics Progress closed chain activities Increase functional activities
Precautions	• None
Suggested Therapeutic Exercises	 Wall sits, shuttle, mini-squats Begin unilateral stance activities, balance training
Progression Criteria	Normal gait mechanics

PHASE IV (12 WEEKS – 6 MONTHS) DATES:

Appointments	PT 1x week weaning down to home workouts
Rehabilitation Goals	 Maximize core/glutes, pelvic stability work Light impact running on Treadmill at 12 weeks progress to plyos and outside funning at 18 weeks - 6 months Gradual return to full unrestricted activities
Precautions	 Post-activity soreness should resolve within 24 hours Avoid knee pain with impact
Suggested Therapeutic Exercises	 Advance Phase III exercises Core/glute, pelvic stability work, eccentric hamstrings, sport/work specific balance and proprioceptive drills
Cardiovascular Exercises	May begin elliptical, bike, and pool as tolerated
Progression Criteria	Physician approval

PHASE V (6 MONTHS – 12 MONTHS) DATES:

Appointments	Schedule MD visit linked to Return to sport testing in PT
Rehabilitation Goals	 Advance functional activity Return to sport-specific activity
Precautions	MD clearance for impact and sport-specific activity after 6 months
Suggested Therapeutic Exercises	Return to sport-specific activity and testing

(Resources used: Dr. Brian Cole, Marc Sherry, PT, DPT, LAT, CSCS(msherry@uwhealth.org) and the UW Health Sports Medicine physician group)

MAMMOTH ORTHOPEDIC INSTITUTE

85 Sierra Park Road • Mammoth Lakes, CA 93546 • 760.924.4084 162 South Main Street • Bishop, CA 93514• 760.872.7766 SIERRA PARK PHYSICAL AND OCCUPATIONAL THERAPY