



Mammoth Orthopedic
Institute

Mammoth Orthopedic Institute ▪ 85 Sierra Park Road ▪ Mammoth Lakes, CA 93546 ▪ 760.924.4084

www.mammothortho.com

ANKLE ARTHROSCOPY with Lateral Ligament Repair

ACTIVITY

- Crutches may help you balance for the first few days; however, you should be non-weight bearing until allowed to do so by orthopedics. Approximately 2 weeks
- You may bend and straighten your knee as much as you like.
- Do not engage in prolonged periods of standing or walking over the first 7-10 days following surgery. Keep the ankle elevated above heart level as much as you are able
- Avoid long periods of sitting (without leg elevated) or long distance traveling for 2 weeks.

DRESSINGS & INCISIONS

- You will be in a well-padded splint following surgery. You should keep this on as placed until your first post-operative visit.
- Please keep the dressing clean and dry; if you are going to shower/bathe, you must protect the dressing with a cast cover. You may not soak in a pool, lake, hot tub, or the ocean until at least 1 week after the sutures have been removed if cleared to do so

PAIN & INFLAMMATION

- Ice - Apply an ice bag wrapped in a dry towel several times per day for 20 minutes for the first week and then as needed for pain relief and inflammation.
- Elevation - Keep your foot elevated above your heart as much as possible for the first 3 to 4 days.
- Pain Medication
 - You have been given a prescription for pain control; please take as directed.
 - If you think you will require a refill on your medication, you **MUST** do so during our regular weekday office hours.
 - If you need additional pain medication you may take Tylenol 500-650 mg every 4-6 hours. Do not take more than 3 grams or 3000 mg in a 24-hour period!
 - Common side effects of the pain medication are:
 - NAUSEA: To decrease nausea, take these medications with food.
 - DROWSINESS: Do not drive a car or operate machinery.
 - ITCHING: You may take Benadryl to alleviate any itching.
 - CONSTIPATION: To decrease constipation, use the stool softener provided (Docusate 250mg) or over-the-counter remedies (Milk of Magnesia, Miralax, etc). Also avoid bananas, rice, apples, toast, or yogurt...as these foods can make you constipated. Getting up and moving around also helps with constipation by “waking up” your intestinal tract.

EMERGENCIES

Please have someone stay with you for the first 24 hours after surgery

Please call the clinic or the orthopedist on-call if:

Drainage soaks the dressings, expands, is foul-smelling, or your incisions are red, warm, and extremely painful

You develop a fever (>101.5°) or chills

You experience leg or calf pain, leg swelling, or difficulty breathing

FOLLOW-UP CARE

Please schedule a follow-up visit for suture removal, and to review your surgery 10-14 days postoperatively.

IF YOU HAVE ANY QUESTIONS OR CONCERNS, PLEASE CALL

Mammoth 760-924-4084

Bishop 760-872-7766

EXERCISES –

When you are comfortable and ready you may perform each exercise 2-3 times a day; it may help to take pain medication 20-30 minutes prior to the exercises and to apply ice after the exercises

Flexion:

Sit in a chair

Place your unoperated leg (B) under the foot of your operated leg (A)

Gently allow the knee to bend with support from your unoperated leg (B)

When you reach your maximum bend, hold for 5 seconds

Perform 10-20 times in a row

Goal = 90° of flexion (bending) by 2 weeks after surgery

Quadriceps Contractions:

Sit or lie on the floor with your operated leg straight

Place a towel roll under the knee

Tighten your thigh and hamstring muscles, causing you to press your knee downward into the towel roll

Hold this position for 10 seconds

Relax your thigh and hamstring muscles

Perform 2-3 sets of 10

Straight Leg Raises:

Lie on the floor

Perform a quadriceps contraction (as stated in the above exercise)

Raise your foot about 6-12" off the floor

Slowly lower your leg back to the floor

Relax your thigh muscle

Perform 2-3 sets of 10