



# *Post Operative Instructions: Tibia and Fibula ORIF/ Nail*

---

## **ACTIVITY**

- Do not bear weight on the operative leg until permitted by your surgeon. Please use crutches to assist with walking.
- Do not engage in prolonged periods of standing or walking the first 7-10 days following surgery.
- Avoid long periods of sitting (without leg elevated) or long distance traveling for 2 weeks.
- You may move your hip and knee as much as you would like.
- You should stretch and move your ankle and foot as much as possible for the first few weeks after surgery.

---

## **BOOT**

- Please keep the boot clean and dry.
- Please wait until the 3rd day after surgery to shower. Remove the boot and just let water run over the incision.
- Do not soak in a tub or pool for a week after the staples are removed. We recommend a sponge bath initially, particularly because it is hard to stand on one leg in the shower, especially if you are on opioid pain medication.

---

## **PAIN & INFLAMMATION**

### Ice

- Apply ice wrapped in a dry towel 20 minutes at a time, several times per day for the first week, then as needed.

### Compression

- Use an ACE wrap or white stocking to reduce swelling.
- The white stocking should be worn for 5–7 days to help prevent blood clots and reduce knee swelling.

### Elevation

- Keep your foot elevated above heart level as much as possible for the first 3–4 days.
- Place a pillow under your calf or foot, not under the knee.

### Pain Medications

- Take your prescribed medications as directed.

#### Tylenol (Acetaminophen)

- First-line for pain.
- Take 1000 mg every 6–8 hours.
- Do not exceed 1000 mg per dose or 4000 mg per day (limit to 3000 mg/day if you have liver issues).
- Do not drink alcohol while taking Tylenol.

#### Oxycodone

- For moderate to severe breakthrough pain only.

- Use sparingly and wean off as soon as possible.

Common side effects include:

- Nausea: Take with food.
- Drowsiness: Do not drive or operate machinery.
- Itching: Benadryl may help.
- Constipation:
  - Use stool softeners (e.g., Senna-Docusate) or OTC options (Mineral Oil, Milk of Magnesia, etc).
  - Avoid bananas, rice, apples, toast, and yogurt (may worsen constipation).
  - Light walking helps stimulate bowel function.

Refills:

- If you think you will need a refill, you must request it during regular weekday office hours.
- 

### **DVT PROPHYLAXIS**

- Take Aspirin 162 mg daily as prescribed.
    - This is not for pain—it's to reduce the risk of blood clots (DVT/pulmonary embolism).
    - If you are at higher risk, you may be placed on Lovenox instead. If you are already on anticoagulation, you may have a different plan.
- 

### **EMERGENCIES**

You must have someone stay with you during the first 48 hours after surgery.

Call the clinic or the on-call orthopedist if:

- Drainage soaks the dressing, increases, smells foul, or if incisions become red, warm, or very painful.
  - You develop a fever >101.5°F or chills.
  - You experience leg or calf pain, swelling, or difficulty breathing.
- 

### **FOLLOW-UP CARE**

- Schedule a follow-up appointment at ~2 weeks postoperatively for a wound check, suture removal, and review of your surgery (if not already scheduled).
-